

# Chloroquine and Hydroxychloroquine Retinal Toxicity Consideration in the Treatment of COVID-19

Paisan Ruamviboonsuk, MD\*, Timothy Y. Y. Lai, MD, FRCOphth†, Andrew Chang, PhD, FRANZCO‡, Chi-Chun Lai, MD§, William F Mieler, MD¶, and Dennis S. C. Lam, MD||\*\*,  
for Asia-Pacific Vitreo-Retina Society

**Abstract:** The proposed doses of chloroquine (CQ) and hydroxychloroquine (HCQ) for treatment of COVID-19 (1000 mg/day for 10 days, CQ; 800 mg first day then 400 mg/day for 5 days, HCQ) in many guidelines worldwide, are considerably higher than the maximum recommended daily safe doses of both agents ( $\leq 2.3$  mg/kg/day, CQ;  $\leq 5.0$  mg/kg/day, HCQ) for development of retinal toxicity. Irreversible retinal damage can occur if the exposure to the safe doses is  $>5$  years. It is not known whether exposure to high doses over a short period of time can also cause the damage. We recommend that before prescribing CQ or HCQ, history of ocular disease should be obtained to avoid the prescription if appropriate. If either agent is to be used, routine baseline ocular examination is not absolutely necessary. Patients who do not have ocular disease should also be informed about the potential risk of retinal toxicity. Both agents, however, have not yet been proven to be beneficial to COVID-19.

**Key Words:** chloroquine, coronavirus, hydroxychloroquine, maculopathy, retinopathy

(Asia Pac J Ophthalmol (Phila) 2020;9:85–87)

With the occurrence of pandemic of the coronavirus disease 2019 (COVID-19) announced by the World Health Organization in early March 2020 and the number of cases still on the rise in all continents in late March, many therapeutic options have been proposed for this novel and potentially fatal disease. Apart from antiviral agents, chloroquine (CQ) and hydroxychloroquine (HCQ) have been examined for their roles in treatment of COVID-19.<sup>1</sup> This may be because both CQ and HCQ have been

postulated to reduce viral replication in other coronavirus infections.<sup>2,3</sup> According to a recent systematic review, there are almost 20 ongoing randomized controlled clinical trials on both medications for treatment of COVID-19 and all of them are in China. The details including dosing regimens of CQ and HCQ in these trials have been summarized in the review.<sup>4</sup> Although there has not yet been a completed clinical trial and the world is waiting eagerly for the results of these trials and other trials on other treatment options, many authorities have chosen to adopt CQ and HCQ in the guidelines for treatment of COVID-19 based on in vitro studies,<sup>2,3</sup> nonrandomized trial,<sup>5</sup> and anecdotal evidence.<sup>6–11</sup> As the therapeutic doses of CQ and HCQ recommended in the trials and guidelines are relatively high compared with the maximum daily safe dose that is related to CQ and HCQ retinal toxicity, this issue of retinal toxicity should be taken into consideration when employing these 2 medications for treatment of COVID-19 worldwide.

According to the recommendation by the American Academy of Ophthalmology, the most significant major risk factors for CQ and HCQ retinal toxicity are high dose and long duration of use.<sup>12</sup> Other risk factors include concomitant renal disease and use of tamoxifen.<sup>12</sup> The maximum daily dose from this recommendation is  $\leq 5.0$  mg/kg real body weight for HCQ, and  $\leq 2.3$  mg/kg real body weight for CQ. As shown in Table 1, the doses of CQ or HCQ for treatment of COVID-19 in various treatment guidelines worldwide are well beyond these recommended dosing regimens.

The Royal College of Ophthalmologists (RCO) in the UK also addressed the importance of safe dose and duration of prescription of CQ and HCQ for the development of retinal toxicity. Although no absolute safe dose was identified, the RCO recommends the daily dose of HCQ to be  $<5$  mg/kg/day for  $<5$  years as relatively safe for retinal toxicity. However, no safe dose of CQ was recommended and the RCO identified those who receive CQ for  $>1$  year as having risk of retinal toxicity.<sup>13</sup>

Despite the fact that the daily doses of both CQ and HCQ for the treatment of COVID-19 exceed the daily safe doses of both agents, the treatment may still be considered relatively safe for retinal toxicity. The toxicity, which causes irreversible retinal damage and visual loss despite ceasing prescription, requires exposure to the safe dose for a long period of time, generally in excess of 5 years. In general, the recommendation for screening for retinal toxicity from CQ and HCQ is within the first year of use as baseline and then annual screening after a year of use for CQ<sup>13</sup> and 5 years of use for HCQ.<sup>12</sup> Both American Academy of Ophthalmology and RCO recommended the screening should be conducted sooner if the major risk factors are present.<sup>12,13</sup> In the case of treatment of COVID-19 using CQ and HCQ, the

From the \*Department of Ophthalmology, Rajavithi Hospital, College of Medicine, Rangsit University, Bangkok, Thailand; †Department of Ophthalmology and Visual Sciences, The Chinese University of Hong Kong, Hong Kong, China; ‡Sydney Eye Hospital, Sydney University, Sydney, NSW, Australia; §Department of Ophthalmology, Chang Gung Memorial Hospital, Linkou Medical Center, Taoyuan, Taiwan; ¶Department of Ophthalmology and Visual Sciences, University of Illinois at Chicago, Chicago, IL, USA; ||International Eye Research Institute of the Chinese University of Hong Kong (Shenzhen), Shenzhen, China; and \*\*C-MER Dennis Lam Eye Centre, C-MER International Eye Care Group Limited, Hong Kong SAR, China.

Submitted February 9, 2020; accepted March 31, 2020.

The authors have no conflicts of interest to declare.

Correspondence: Paisan Ruamviboonsuk, Department of Ophthalmology, Rajavithi Hospital, College of Medicine, Rangsit University, Bangkok, Thailand 10400. E-mail: paisan.tr.s@gmail.com, Timothy Lai, M.D., FRCOphth, Department of Ophthalmology and Visual Sciences, The Chinese University of Hong Kong, Hong Kong, China 999077. E-mail: tyylai@cuhk.edu.hk.

Copyright © 2020 Asia-Pacific Academy of Ophthalmology. Published by Wolters Kluwer Health, Inc. on behalf of the Asia-Pacific Academy of Ophthalmology. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ISSN: 2162-0989

DOI: 10.1097/APO.0000000000000289

**Table 1.** Comparison Between the Maximum Daily Safe Doses of Chloroquine and Hydroxychloroquine for Development of Retinal Toxicity and the Recommended Doses in COVID-19 Treatment Guidelines From Different Countries

	Maximum Recommended Daily Safe Dose for Retinal Toxicity	Dosing Regimen in Dutch Guideline <sup>7</sup>	Dosing Regimen in Belgian Guideline <sup>9</sup>	Dosing Regimen in Italian Guideline <sup>11</sup>	Dosing Regimen in Chinese Guideline <sup>8</sup>	Dosing Regimen in Thai Guideline <sup>6</sup>
Chloroquine	≤2.3 mg/kg/day Estimated global dose* = 142.6 mg/day	900mg Day 1, then 600 mg/day up to 5 days	900 mg Day 1, then 600 mg/day up to 5 days	1000 mg/day for 10 days	1000 mg/day for 10 days	500 mg/day for at least 5 days
Hydroxychloroquine	≤5.0 mg/kg/day Estimated global dose* = 310 mg/day	800mg Day 1, then 400 mg/day up to 5 days	800 mg Day 1, then 400 mg/day up to 5 days	800 mg Day 1, then 400 mg/day for 10 days	N/A	800 mg Day 1, then 400 mg/day for at least 5 days

\*The estimated global doses are estimated according to the global average weight of adults (62 kg).<sup>17</sup>

major risk factor is the use of higher than generally recommended dosage, although over a relatively short period of time, that is for about a week. There has not yet been a report on retinal toxicity associated with this kind of treatment. Nonetheless, it has been reported that retinal toxicity can develop even after <1 year of high dose of HCQ use (1000 mg daily) in an oncology trial.<sup>14</sup>

In this report, 2 of 7 patients who received the high dose of HCQ showed abnormalities of the macula on retinal imaging modalities and multifocal electroretinogram without visual symptoms. These patients did not have any known risk factors, such as renal disease, concomitant retinotoxic agents, or co-existing retinal disease. It is not known whether the retinal toxicity from high-dose CQ and HCQ is underreported in the literature due to suboptimal and nonuniform ocular screening methods.<sup>14</sup>

Both CQ and HCQ are known for their binding affinity with melanin in retinal pigment epithelium which can be a mechanism of the toxic effects. Both agents have also been shown to cause damage to the photoreceptor layer and outer nuclear layer of the retina, whereas CQ can cause damage to the inner retina as well. Light absorption and metabolism of cone cells may also play roles for the damages. These mechanisms may lead to clinically characteristic “bull’s eye” maculopathy after chronic exposure to both agents even in the safe dose.<sup>15</sup> It is not known whether exposure to the high dose over a short period may also cause similar cellular damages as with the chronic exposure. Given that patients with COVID-19 who may require treatment are commonly older patients, it is possible that some may already have coexistent age-related macular degeneration. It is still not known whether the diseased macula would be more vulnerable to damage with exposure to the high dose of either CQ or HCQ even over a short duration.

Routine baseline ocular examination is not absolutely necessary for patients with COVID-19 who are undergoing treatment with CQ and HCQ but should be considered if manpower and expertise are available and extreme precautions should be taken during the examination. It is relevant, however, to take a history of ocular disease, particularly macular disease, in patients with COVID-19 who are older than 50 years before prescribing CQ or HCQ as treatment, to rule out age-related macular degeneration or other macular abnormalities. Coexistent retinal pathology is listed as a contraindication of using CQ and HCQ in patients with COVID-19 in the treatment guideline of Belgium.<sup>9</sup> As treatment with CQ or HCQ is not yet proven to be beneficial, but instead can be harmful,<sup>16</sup> in COVID-19, choosing other options of treatment in this group of patients with the preexisting disease may be more appropriate. For patients in whom CQ or HCQ is still considered as a treatment option, the potential benefits and risks of retinal toxicity and other systemic complications shall be thoroughly discussed with patients and well documented on written consent form before the treatment or trial of CQ or HCQ. Following recovery from COVID-19 with the treatment using CQ or HCQ, the patients should also be informed to visit ophthalmologists if they encounter any abnormal visual symptoms.

In summary, the bottom line at the present time is that neither CQ nor HCQ has been proven to be effective in the treatment of COVID-19, although there is certainly a vast interest in its possible benefit. Further controlled clinical trial data will be necessary to help better address this issue. Despite the current situation of COVID-19 pandemic, many adverse effects of either CQ or HCQ should still be weighed against its potential benefit. For retinal toxicity, the risk of having irreversible retinal damage

and visual loss may outweigh the unproven benefit of both agents in some patients. Detecting the risk is easy. It can be done by simply taking a history of previous or co-existing ocular disease from the patients, then other options of treatment should be considered if appropriate.

### ACKNOWLEDGEMENT

The authors acknowledge Varis Ruamviboonsuk, MD, in assisting manuscript preparation.

### REFERENCES

- Zhou D, Dai SM, Tong Q. COVID-19: a recommendation to examine the effect of hydroxychloroquine in preventing infection and progression. *J Antimicrob Chemother*. 2020; doi: 10.1093/jac/dkaa114.
- Yao X, Ye F, Zhang M, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis*. 2020; doi: 10.1093/cid/ciaa237.
- Liu J, Cao R, Xu M, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov*. 2020;6:16.
- Cortegiani A, Ingoglia G, Ippolito M, Giarratano A, Einav S. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care*. 2020; doi: 10.1016/j.jcrc.2020.03.005.
- Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents*. 2020. doi: 10.1016/j.ijantimicag.2020.105949.
- Department of Medical Services Ministry of Public Health Thailand. Thai Treatment Guideline for COVID-19. 2020. Available at: [https://ddc.moph.go.th/viralpneumonia/file/g\\_health\\_care/G2\\_new3\\_1.pdf](https://ddc.moph.go.th/viralpneumonia/file/g_health_care/G2_new3_1.pdf). Accessed March 26, 2020.
- Vollaard A, Gieling E, Linden P van der, et al. Medicamenteuze behandelopties bij patiënten met COVID-19 (infecties met SARS-CoV-2). 2020. Available at: <https://swab.nl/nl/covid-19>. Accessed March 26, 2020.
- Multicenter collaboration group of Department of Science and Technology of Guangdong Province and Health Commission of Guangdong Province for chloroquine in the treatment of novel coronavirus pneumonia. [Expert consensus on chloroquine phosphate for the treatment of novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43:185–188.
- Institute of Tropical Medicine Antwerp, Universiteit Antwerpen, CHU Saint-Pierre, et al. Interim Clinical Guidance for Patients Suspected of/ Confirmed With Covid-19 in Belgium. 2020: 1–13. Available at: [https://epidemiology.wiv-isp.be/ID/Documents/Covid19/COVID-19\\_InterimGuidelines\\_Treatment\\_ENG.pdf](https://epidemiology.wiv-isp.be/ID/Documents/Covid19/COVID-19_InterimGuidelines_Treatment_ENG.pdf). Accessed March 23, 2020.
- Centers for Disease Control and Prevention. Therapeutic Options for COVID-19 Patients | CDC. 2020. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/therapeutic-options.html>. Accessed March 23, 2020.
- Nicastri E, Petrosillo N, Bartoli TA, et al. National Institute for the Infectious Diseases “L. Spallanzani” IRCCS. Recommendations for COVID-19 Clinical Management. *Infect Dis Rep*. 2020; doi: 10.4081/idr.2020.8543.
- Marmor MF, Kellner U, Lai TY, Melles RB, Mieler WF, Lum F. Recommendations on screening for chloroquine and hydroxychloroquine retinopathy (2016 revision). *Ophthalmology*. 2016;123:1386–1394.
- Yusuf IH, Foot B, Galloway J, et al. The Royal College of Ophthalmologists recommendations on screening for hydroxychloroquine and chloroquine users in the United Kingdom: executive summary consensus-statement. *Eye (Lond)*. 2018;32:1168–1173.
- Leung LS, Neal JW, Wakelee HA, Sequist LV, Marmor MF. Rapid onset of retinal toxicity from high-dose hydroxychloroquine given for cancer therapy. *Am J Ophthalmol*. 2015;160:799–805.
- Yusuf IH, Sharma S, Luqmani R, Downes SM. Hydroxychloroquine retinopathy. *Eye (Lond)*. 2017;31:828–845.
- Guastalegname M, Vallone A. Could chloroquine /hydroxychloroquine be harmful in Coronavirus Disease 2019 (COVID-19) treatment? *Clin Infect Dis*. 2020; doi: 10.1093/cid/ciaa321.
- Walpole SC, Prieto-Merino D, Edwards P, Cleland J, Stevens G, Roberts I. The weight of nations: an estimation of adult human biomass. *BMC Public Health*. 2012;12:439.